Brain Research, 523 (1990) 359–361
Elsevier

Author Index

Adeymo, O.M., see Shapira, S., 291
Adrover, E., see Gallego, R., 298
Agarwal, S.K. and Calaresu, F.R., Reciprocal connections between nucleus tractus solitarii and rostral ventrolateral medulla, 305
Akiyama, H., see Tooyama, I., 273
Alberch, J., see Arenas, E., 51
Albin, R.L., see Makowiec, R.L., 309
Alexander, N., Kaneda, N., Ishii, A., Mogi, M., Harada, M. and Nagatsu, T., Right–left asymmetry of tyrosine hydroxylase in rat median eminence: influence of arterial baroreflex nerves, 195
Allen, S.L., see Myall, O.T., 295
Amano, S., Obata, T., Hazama, F., Kashiro, N. and Shimada, M., Hypoxia prevents seizures and neuronal damages of the hippocampus induced by kainic acid in rats, 121
Amundson, R., see Barron, K.D., 219
Appenteng, K., see Moore, J.A., 139
Arenas, E., Alberch, J., Sanchez Arroyos, R. and Marsal, J., Effect of opioids on acetylcholine release evoked by K^+ or glutamic acid from rat neostriatal slices, 51
Arias, P., Szwarcfarb, B., De Rondina, D.C., Carbone, S., Sverdlik, R. and Moguilevsky, J.A., In vivo and in vitro studies of the effect of the serotoninergic system on luteinizing hormone and luteinizing hormone-releasing hormone secretion in prepubertal and peripubertal female rats, 57
Arrang, J.-M., see Lin, J.-S., 325

Ballanyi, K., see Dörner, R., 111
Barclay, S.R. and Harding, C.F., Differential modulation of monoamine levels and turnover rates by estrogen and/or androgen in hypothalamic and vocal control nuclei of male zebra finches, 251
Barron, K.D., Marciano, F.F., Amundson, R. and Mankes, R., Perineuronal glial responses after axotomy of central and peripheral axons. A comparison, 219
Baudry, M., see Jensen, J., 313
Beinfeld, M.C., see Eiden, L.E., 245
Bermúdez-Rattoni, F., see López-García, J.C., 100
Bermúdez-Rattoni, F., see López-García, J.C., 105
Bermúdez-Rattoni, F., see Paredes, R., 331
Bernard, J.F., Huang, G.F. and Besson, J.M., Effect of noxious somesthetic stimulation on the activity of neurons of the nucleus centralis of the amygdala, 347
Besson, J.M., see Bernard, J.F., 347
Bongianni, F., Christenson, J., Hökfelt, T. and Grillner, S., Neuropeptide Y-immunoreactive spinal neurons make close appositions on axons of primary sensory afferents, 337
Boyajian, C.L., see Cooper, D.M.F., 143
Brachet, P., see Chakrabarti, S., 11
Braschler, U., see Remé, C.E., 167
Butcher, L.L., see Di Patre, P.L., 316

Calaresu, F.R., see Agarwal, S.K., 305
Cano, J., see Vizuete, M.L., 281
Carbone, S., see Arias, P., 57
Cha, J.-H.J., see Makowiec, R.L., 309
Chakrabarti, S., Sima, A.A.F., Lee, J., Brachet, P. and Dicou, E., Nerve growth factor (NGF), proNGF and NGF receptor-like immunoreactivity in BB rat retina, 11
Chase, T.N., see Weick, B.G., 16
Chau, R.M.W., see Jen, L.S., 156
Chen, W.Z., see Jen, L.S., 156
Chowdrey, H.S., see Jessop, D.S., 1
Christenson, J., see Bongianni, F., 337
Colpaert, F.C., see Lategan, A.J., 134
Colwell, C.S., Ralph, M.R. and Menaker, M., Do NMDA receptors mediate the effects of light on circadian behavior?, 117
Cooper, D.M.F., Boyajian, C.L., Goldsmith, P.K., Unson, C.G. and Spiegel, A., Differential expression of low molecular weight form of G_s-α in neostriatum and cerebellum: correlation with expression of calmodulin-independent adenylyl cyclase, 143
Crawley, J.N., see Laitinen, K., 342

Damez-Kinselle, I., see Guastavino, J.-M., 199
Depaulis, A., see Vergnes, M., 87
De Rondina, D.C., see Arias, P., 57
De Witte, Ph.D., see Laitinen, K., 342
Dicou, E., see Chakrabarti, S., 11
Di Patre, P.L., Oh, J.D., Simmons, J.M., and Butcher, L.L., Intrafimbrial colchicine produces transient impairment of radial-arm maze performance correlated with morphologic abnormalities of septohippocampal neurons expressing cholinergic markers and nerve growth factor receptor, 316
Dörner, R., Ballanyi, K. and Schlue, W.-R., Glutaminergic responses of neuropile glial cells and Retzius neurones in the leech central nervous system, 111
Drost, D., see Prato, F.S., 301

Eiden, L.E., Mezey, E., Eskay, R.L., Beinfeld, M.C. and Palkovits, M., Neuropeptide content and connectivity of the rat claustrum, 245
El-Yassir, N. and Fleetwood-Walker, S.M., A 5-HT_1-type receptor mediates the antinociceptive effect of nucleus raphe magnus stimulation in the rat, 92
Engber, T.M., see Weick, B.G., 16
Eskay, R.L., see Eiden, L.E., 245
Eugenin, J., Larrain, C. and Zapata, P., Functional recovery of the ventilatory chemoreflexes after partial chronic denervation of the nucleus tractus solitarius, 263

Favard, C., Simon, A., Vigny, A. and Nguyen-Legros, J., Ultrastructural evidence for a close relationship between dopamine cell processes and blood capillary walls in *Macaca* monkey and rat retina, 127
Fernández-Ruiz, J., see López-García, J.C., 105
Fernández-Ruiz, J., see Paredes, R., 331
Feuerstein, G., see Shapira, S., 291
Fibiger, H.C., see Robertson, G.S., 288
Fleetwood-Walker, S.M., see El-Yassir, N., 92
Frappier, J.R.H., see Prato, F.S., 301
French, A.S., see Stockbridge, L.L., 161

Gallego, R. and Adrover, E., Axotomy decreases conduction velocity of unmyelinated sensory fibres, 298
Garbarg, M., see Lin, J.-S., 325
Gibo, D.M., see Zagon, I.S., 62
Gilman, S., see Makowiec, R.L., 309
Ginzburg, R. and Seltzer, Z., Subarachnoid spinal cord transplantation of adrenal medulla suppresses chronic neuropathic pain behavior in rats, 147
Girman, S.V. and Golovina, I.L., Electrophysiological properties of embryonic neocortex transplants replacing the primary visual cortex of adult rats, 78
Goldsmith, P.K., see Cooper, D.M.F., 143
Golovina, I.L., see Girman, S.V., 78
Grillner, S., see Bongianni, F., 337
Gruenstein, E.I., see Lomneth, R., 69
Guastavino, J.-M., Sotelo, C. and Damez-Kinselle, I., *Hot-foot* murine mutation: behavioral effects and neuroanatomical alterations, 199

Han, J.-S., see Wang, X.-J., 5
Harada, M., see Alexander, N., 195
Harding, C.F., see Barclay, S.R., 251
Hazama, F., see Amano, S., 121
Hearns, M., see Luine, V., 321
Herrera, A.J., see Vizuete, M.L., 281
Hökfelt, T., see Bongianni, F., 337
Huang, G.F., see Bernard, J.F., 347

Ishii, A., see Alexander, N., 195
Ito, Y., see Masuda, T., 356

Jen, L.S., Tsang, D., Chau, R.M.W. and Chen, W.Z., The cholinergic system in retinal transplants in rats, 156
Jensen, J., Lynch, G. and Baudry, M., Regional differences in the activation of synaptosomal mitochondrial Ca^{2+} uptake by spermine in rat brain, 313
Jepson-Innes, K., see Zill, S.N., 211
Jessop, D.S., Chowdrey, H.S. and Light-

man, S.L., Inhibition of rat corticotropin-releasing factor and adrenocorticotropin secretion by an osmotic stimulus, 1
Jouvet, M., see Lin, J.-S., 325

Kadar, T., see Shapira, S., 291
Kaneda, N., see Alexander, N., 195
Kashiro, N., see Amano, S., 121
Kavaliers, M., see Prato, F.S., 301
Kimura, H., see Tooyama, I., 273
Krauthamer, G.M., see Yamasaki, D.S.G., 188
Krug, M., see Schuster, T., 171

Laitinen, K., Crawley, J.N., Mefford, I.N. and De Witte, Ph.D., Neurotensin and cholecystokinin microinjected into the ventral tegmental area modulate microdialysate concentrations of dopamine and metabolites in the posterior nucleus accumbens, 342
Larrain, C., see Eugenin, J., 263
Lategan, A.J., Marien, M.R. and Colpaert, F.C., Effects of locus coeruleus lesions on the release of endogenous dopamine in the rat nucleus accumbens and caudate nucleus as determined by intracerebral microdialysis, 134
Lee, J., see Chakrabarti, S., 11
Lee, T.Y., see Prato, F.S., 301
Levenson, D. and Rosenbluth, J., Electrophysiologic changes accompanying Wallerian degeneration in frog sciatic nerve, 230
Lightman, S.L., see Jessop, D.S., 1
Lin, J.-S., Sakai, K., Vanni-Mercier, G., Arrang, J.-M., Garbarg, M., Schwartz, J.-C. and Jouvet, M., Involvement of histaminergic neurons in arousal mechanisms demonstrated with H_3-receptor ligands in the cat, 325
Little, H.J., see Whittington, M.A., 237
Lomneth, R., Medrano, S. and Gruenstein, E.I., The role of transmembrane pH gradients in the lactic acid induced swelling of astrocytes, 69
López-García, J.C., Bermúdez-Rattoni, F. and Tapia, R., Release of acetylcholine, γ-aminobutyrate, dopamine and glutamate, and activity of some related enzymes, in rat gustatory neocortex, 100
López-García, J.C., Fernández-Ruiz, J., Bermúdez-Rattoni, F. and Tapia, R., Correlation between acetylcholine release and recovery of conditioned taste aversion induced by fetal neocortex grafts, 105
Luine, V. and Hearns, M., Spatial memory deficits in aged rats: contributions of the cholinergic system assessed by ChAT, 321
Lynch, G., see Jensen, J., 313

Machado, A., see Vizuete, M.L., 281
Makowiec, R.L., Albin, R.L., Cha, J.-H.J., Young, A.B. and Gilman, S., Two types of quisqualate receptors are decreased in human olivopontocerebellar atrophy cerebellar cortex, 309
Mankes, R., see Barron, K.D., 219
Marciano, F.F., see Barron, K.D., 219
Marescaux, C., see Vergnes, M., 87
Marien, M.R., see Lategan, A.J., 134
Marsal, J., see Arenas, E., 51
Masuda, T., Yamazaki, Y. and Ito, Y., Comparison between extracellular and tissue fluctuation of dopamine and related metabolites in striatum under hypoxia, 356
Medrano, S., see Lomneth, R., 69
Mefford, I.N., see Laitinen, K., 342
McGeer, P.L., see Tooyama, I., 273
McLaughlin, P.J., see Zagon, I.S., 62
McLean, W.G., see Myall, O.T., 295
Menaker, M., see Colwell, C.S., 117
Mezey, E., see Eiden, L.E., 245
Mogi, M., see Alexander, N., 195
Moguilevsky, J.A., see Arias, P., 57
Moore, J.A. and Appenteng, K., Contrasting effects of urethane and pentobarbitone anaesthesia on the electrical properties of rat jaw-elevator motoneurones, 139
Munz, K., see Remé, C.E., 167
Myall, O.T., Allen, S.L. and McLean, W.G., The effect of acrylamide on the induction of ornithine decarboxylase in the dorsal root ganglion of the rat, 295

Nagatsu, T., see Alexander, N., 195
Nguyen-Legros, J., see Favard, C., 127

Obata, T., see Amano, S., 121
O'Connor, D.T., see Powers, R.E., 151
Oh, J.D., see Di Patre, P.L., 316
Otani, K., see Shigenaga, Y., 23

Palkovits, M., see Eiden, L.E., 245
Paredes, R.G., Piña, A.L., Fernández-Ruiz, J. and Bermúdez-Rattoni, F., Fetal brain transplants induce recovery of male sexual behavior in medial preoptic area-lesioned rats, 331
Piña, A.L., see Paredes, R., 331
Powers, R.E., O'Connor, D.T. and Price, D.L., Noradrenergic innervation of human inferior olivary complex, 151
Prato, F.S., Frappier, J.R.H., Shivers, R.R., Kavaliers, M., Zabel, P., Drost, D. and Lee, T.Y., Magnetic resonance imaging increases the blood–brain barrier permeability to 153-gadolinium diethylenetriaminepentaacetic acid in rats, 301
Price, D.L., see Powers, R.E., 151

Ralph, M.R., see Colwell, C.S., 117
Remé, C.E., Braschler, U., Wirz-Justice, A. and Munz, K., Disk-shedding in the rat retina: lithium dampens the circadian rhythm but potentiates the light response, 167
Robertson, G.S., Vincent, S.R., and Fibiger, H.C., Striatonigral projection neurons contain D_1 dopamine receptor-activated c-*fos*, 288
Rosenbluth, J., see Levenson, D., 230
Rustioni, A., see Weinberg, R.J., 351

Sakai, K., see Lin, J.-S., 325
Sanchez, Arroyos, R., see Arenas, E., 51
Sanders, E.J., see Stockbridge, L.L., 161
Santiago, M., see Vizuete, M.L., 281
Schlue, W.-R., see Dörner, R., 111
Schuster, T., Krug, M. and Wenzel, J., Spinules in axospinous synapses of the rat dentate gyrus: changes in density following long-term potentiation, 171
Schwartz, J.-C., see Lin, J.-S., 325
Seltzer, Z., see Ginzburg, R., 147
Shapira, S., Kadar, T., Adeymo, O.M. and Feuerstein, G., Selective hippocampal lesion following ω-conotoxin administration in rats, 291
Shigenaga, Y., Otani, K. and Suemune, S., Morphology of central terminations of low-threshold trigeminal primary afferents from facial skin in the cat — intra-axonal staining with HRP, 23
Shimada, M., see Amano, S., 121
Shivers, R.R., see Prato, F.S., 301
Sima, A.A.F., see Chakrabarti, S., 11
Simmons, J.M., see Di Patre, P.L., 316
Simon, A., see Favard, C., 127
Sleight, S., see Wehner, J.M., 181
Sotelo, C., see Guastavino, J.-M., 199
Spiegel, A., see Cooper, D.M.F., 143
Stockbridge, L.L., French, A.S. and Sanders, E.J., Dissociation and culture of mechanosensory neurons for patch clamp analysis, 161
Suemune, S., see Shigenaga, Y., 23
Susel, Z., see Weick, B.G., 16
Sverdlik, R., see Arias, P., 57
Szwarcfarb, B., see Arias, P., 57

Tapia, R., see López-García, J.C., 100
Tapia, R., see López-García, J.C., 105
Tooyama, I., Kimura, H., Akiyama, H. and McGeer, P.L., Reactive microglia express class I and class II major histocompatibility complex antigens in Alzheimer's disease, 273
Tracey, D.J., see Weinberg, R.J., 351
Tsang, D., see Jen, L.S., 156

Unson, C.G., see Cooper, D.M.F., 143
Upchurch, M., see Wehner, J.M., 181

Vanni-Mercier, G., see Lin, J.-S., 325
Vergnes, M., Marescaux, C. and Depaulis, A., Mapping of spontaneous spike and wave discharges in Wistar rats with genetic generalized non-convulsive epilepsy, 87
Vigny, A., see Favard, C., 127
Vincent, S.R., see Robertson, G.S., 288
Vizuete, M.L., Herrera, A.J., Santiago, M., Machado, A. and Cano, J., Effects of enucleation on postnatal development of catecholamines and serotonin metabolism in the superior colliculus of the rat, 281

Walters, J.R., see Weick, B.G., 16
Wang, X.-H., see Wang, X.-J., 5
Wang, X.-J., Wang, X.-H. and Han, J.-S., Cholecystokin in octapeptide antagonized opioid analgesia mediated by μ- and κ- but not δ-receptors in the spinal cord of the rat, 5
Wehner, J.M., Sleight, S. and Upchurch, M., Hippocampal protein kinase C activity is reduced in poor spatial learners, 181
Weick, B.G., Engber, T.M., Susel, Z., Chase, T.N. and Walters, J.R., Responses of substantia nigra pars reticulata

neurons to GABA and SKF 38393 in 6-hydroxydopamine-lesioned rats are differentially affected by continuous and intermittent levodopa administration, 16
Weinberg, R.J., Tracey, D.J. and Rustioni, A., Extracellular labeling of unmyelinated dorsal root ganglion after WGA-HRP injections in spinal ganglia, 351
Wenzel, J., see Schuster, T., 171
Whittington, M.A. and Little, H.J., Patterns of changes in field potentials in the isolated hippocampal slice on withdrawal from chronic ethanol treatment of mice in vivo, 237
Wirsig-Wiechmann, C.R., The nervus terminalis in the chick: a FMRFamide-immunoreactive and AChE-positive nerve, 175
Wirz-Justice, A., see Remé, C.E., 167

Yamasaki, D.S.G. and Krauthamer, G.M., Somatosensory neurons projecting from the superior colliculus to the intralaminar thalamus in the rat, 188
Yamazaki, M., see Masuda, T., 356
Young, A.B., see Makowiec, R.L., 309

Zabel, P., see Prato, F.S., 301
Zagon, I.S., Gibo, D.M. and McLaughlin, P.J., Adult and developing human cerebella exhibit different profiles of opioid binding sites, 62
Zapata, P., see Eugenin, J., 263
Zill, S.N. and Jepson-Innes, K., Functions of a proprioceptive sense organ in freely moving insects: characteristics of reflexes elicited by stimulation of the locust metathoracic femoral chordotonal organ, 211

(contents continued)

Neuropeptide Y-immunoreactive spinal neurons make close appositions on axons of primary sensory afferents
F. Bongianni, J. Christenson, T. Hökfelt and S. Grillner (Sweden) 337
Neurotensin and cholecystokinin microinjected into the ventral tegmental area modulate microdialysate concentrations of dopamine and metabolites in the posterior nucleus accumbens
K. Laitinen, J.N. Crawley, I.N. Mefford and Ph. De Witte (U.S.A., Belgium) 342
Effect of noxious somesthetic stimulation on the activity of neurons of the nucleus centralis of the amygdala
J.F. Bernard, G.F. Huang and J.M. Besson (France) 347
Extracellular labeling of unmyelinated dorsal root ganglion after WGA-HRP injections in spinal ganglia
R.J. Weinberg, D.J. Tracey and A. Rustioni (U.S.A.) 351
Comparison between extracellular and tissue fluctuation of dopamine and related metabolites in striatum under hypoxia
T. Masuda, M. Yamazaki and Y. Ito (Japan) 356
Author Index 359